

A NOTE ON THE KOSZUL COMPLEX IN DEFORMATION QUANTIZATION

A. FERRARIO, C. A. ROSSI, AND T. WILLWACHER

ABSTRACT. The aim of this short note is to present a proof of the existence of an A_∞ -quasi-isomorphism between the A_∞ - $S(V^*)$ - $\wedge(V)$ -bimodule K , introduced in [1], and the Koszul complex $K(V)$ of $S(V^*)$, viewed as an A_∞ - $S(V^*)$ - $\wedge(V)$ -bimodule, for V a finite-dimensional (complex or real) vector space.

1. INTRODUCTION

The main result of [1] is a Formality Theorem in presence of two subspaces U_1, U_2 of a real or complex finite-dimensional vector space V , constructed using the graphical techniques of Kontsevich [7]. Such a Formality Theorem has interesting by-products even in the simplest case, when $U_1 = V$ and $U_2 = \{0\}$:

- (1) It implies that the deformation quantization of $A = \mathcal{O}_V = S(V^*)$ [7] and of $B = \mathcal{O}_{V^*[1]} = \wedge(V)$ [4] w.r.t. a given polynomial Poisson structure on V preserves Koszul duality.
- (2) It yields a proof of a conjecture of B. Shoikhet [2, 9] about the possibility of realizing Kontsevich's deformed algebra $A_\hbar = A[[\hbar]]$ [7] w.r.t. a polynomial Poisson structure *via* generators and relations.

In this particular context, the main object appearing in the Formality Theorem is $K = \mathbb{K}$ (here, $\mathbb{K}$ is any field of characteristic 0 containing $\mathbb{R}$), endowed with a nontrivial A_∞ - A - B -bimodule structure. This bimodule structure is constructed explicitly in [1, Subsection 6.2], using Kontsevich's graphical techniques, specializing earlier results of Cattaneo–Felder [3, 4]. The main result of this note is the following

Theorem 1.1. *The A_∞ -bimodule K is A_∞ -quasi-isomorphic to the Koszul complex $K(V)$ of A , endowed with the usual A - B -bimodule structure.*

Theorem 1.1 has been previously stated as a conjecture, see [1, Conjecture 1.3], where the A_∞ - A - B -bimodule has been introduced.

In fact, the Koszul complex, which is a dg A - B -bimodule, has been considered in the framework of Deformation Quantization, in [10], where the author used Tamarkin's Formality in order to quantize both algebras A and B and the Koszul complex itself, in order to prove that Koszul duality is preserved by Deformation Quantization.

The approach of [1] makes use of Kontsevich's Formality: the A_∞ - A - B -bimodule structure originates from perturbative expansion of the Poisson Sigma model, and it turns out that it realizes, in the *A_{infty}*-framework, Koszul duality between A and B : it turns out that it behaves well w.r.t. Deformation Quantization, hence it is the right candidate for showing that Kontsevich's Deformation Quantization techniques preserve also Koszul duality.

For the proof we check that both morphisms in the sequence

$$K(V) \hookrightarrow A \otimes_A K \rightarrow K$$

are quasi-isomorphisms of A_∞ -bimodules. Here $(-\otimes_A -)$ denotes a tensor product of A_∞ -bimodules, defined in Section 3. As for ordinary bimodules, one verifies that the tensor product with the algebra itself, i.e., $(A \otimes_A -)$, is (quasi-isomorphic to) the identity, and hence the right arrow is a quasi-isomorphism. The fact that the left arrow is a quasi-isomorphism of A_∞ -bimodules is due to a peculiarity of the bimodule structure on K , and is proven in Section 5.

Remark 1.2. It has been pointed out to us by B. Keller and A. Khoroshkin independently that Theorem 1.1 should follow from the results of [1] and general arguments of homological algebra: however, we do not know, for the time being, a rigorous proof in this framework.

Acknowledgments. We kindly thank G. Felder and D. Calaque for many inspiring discussions, suggestions and constructive criticism. We also thank B. Keller and A. Khoroshkin for the comments in Remark 1.2, although this paper may not answer to their criticism in the way they intended it. Finally, we thank J. Stasheff and the referee for having carefully read this paper and for many constructive critics and editorial improvements. The second author acknowledges the stimulating working atmosphere of the FIM of the ETH, Zürich, where this work has been accomplished.

2000 *Mathematics Subject Classification.* 16E45, 81R60.

Key words and phrases. Koszul duality; A_∞ -algebras and -bimodules; Deformation Quantization.

2. NOTATION AND CONVENTIONS

Throughout the paper, $\mathbb{K}$ is a field of characteristic 0, which contains $\mathbb{R}$. V is a finite-dimensional vector space over $\mathbb{K}$, V^* its dual. Further, we denote by $\{x_i\}$, $i = 1, \dots, d = \dim V$, a basis of V^* , which yields automatically global linear coordinates on V , which we denote by y_i .

Let $\mathbf{Mod}_{\mathbb{K}}$ be the monoidal category of graded vector spaces, with graded tensor product, and with inner spaces of morphisms (i.e. we consider morphisms, which are finite sums of morphisms of any degree); $[\bullet]$ denotes the degree-shifting functor on $\mathbf{Mod}_{\mathbb{K}}$. In particular, the identity morphism of an object M of $\mathbf{Mod}_{\mathbb{K}}$ induces a canonical isomorphism $s : M \rightarrow M[1]$ of degree -1 with inverse $s^{-1} : M[1] \rightarrow M$ (suspension and de-suspension isomorphisms): for the sake of simplicity, we will use the following short-hand notation

$$(v_1 | \dots | v_n) = s(v_1) \otimes \dots \otimes s(v_n).$$

The degree of an element m of a homogeneous component of an object M of $\mathbf{Mod}_{\mathbb{K}}$ is denoted by $|m|$. Unadorned tensor products are meant to be over $\mathbb{K}$.

A (possibly curved) A_{∞} -algebra A over $\mathbb{K}$ is equivalent to the structure of a codifferential cofree coalgebra with counit on $T(A[1]) = \bigoplus_{n \geq 0} A[1]^{\otimes n}$, for an object A of $\mathbf{Mod}_{\mathbb{K}}$. The codifferential d_A is uniquely determined by its Taylor components

$$d_A^n : A[1]^{\otimes n} \rightarrow A[1], \quad n \geq 0,$$

all of degree 1, and the condition that d_A squares to 0 translates into an infinite family of quadratic relations between its Taylor components. We further set $m_A^n = s^{-1} \circ d_A^n \circ s^{\otimes n}$. By construction, m_A^n are $\mathbb{K}$ -linear maps from $A^{\otimes n}$ to A of degree $2 - n$. We refer to m_A^0 as to the curvature of A . It is an element of A of degree 2, which measures the failure of (A, m_A^1) to be a differential graded (shortly, from now on, dg) vector space over $\mathbb{K}$. If $m_A^0 = 0$, then A is said to be flat.

Finally, given two (possibly curved) A_{∞} -algebras A, B , an A_{∞} - A - B -bimodule structure on an object K of $\mathbf{Mod}_{\mathbb{K}}$ is equivalent to the structure of a codifferential cofree bicomodule on $T(A[1]) \otimes K[1] \otimes T(B[1])$. As for A_{∞} -algebras, such a codifferential d_K is uniquely determined by its Taylor components

$$d_K^{m,n} : A[1]^{\otimes m} \otimes K[1] \otimes B[1]^{\otimes n} \rightarrow K[1], \quad m, n \geq 0,$$

all of degree 1. As before, we introduce the maps $m_K^{m,n} = s^{-1} \circ d_K^{m,n} \circ s^{\otimes m+1+n}$, of degree $1 - m - n$. The condition that d_K squares to 0 is equivalent to an infinite family of quadratic relations between the Taylor components of d_A , d_B and d_K . For more details on A_{∞} -bimodules over (possibly curved) A_{∞} -algebras, we refer to [1, Sections 3 and 4].

Remark 2.1. We observe that, if A and B are both flat, then an A_{∞} - A - B -bimodule structure on K yields both a left A_{∞} - A - and right A_{∞} - B -module structure on K in the sense of [5, 8], but, if either A or B or both are curved, then the A_{∞} -bimodule structure does not restrict to (left or right) A_{∞} -module structures, see e.g. [11] and [1, Subsection 4.1].

3. THE TENSOR PRODUCT OF A_{∞} -BIMODULES

We consider now three (possibly curved) A_{∞} -algebras A, B and C . Furthermore, we consider an A_{∞} - A - B -bimodule K_1 and an A_{∞} - B - C -bimodule K_2 . The tensor product of K_1 and K_2 over B , as an element of $\mathbf{Mod}_{\mathbb{K}}$, is defined as

$$K_1 \underline{\otimes}_B K_2 = K_1 \otimes T(B[1]) \otimes K_2.$$

The A_{∞} -structures on A, B, C, K_1 and K_2 determine a unique structure of A_{∞} - A - C -bimodule over $K_1 \underline{\otimes}_B K_2$, which we now describe explicitly. By the arguments in Section 2, it suffices to construct a codifferential on the cofree bicomodule

$$T(A[1]) \otimes (K_1 \underline{\otimes}_B K_2)[1] \otimes T(C[1]) \cong T(A[1]) \otimes K_1[1] \otimes T(B[1]) \otimes K_2[1] \otimes T(C[1]),$$

where the isomorphism is induced by suspension and de-suspension, and has degree -1 ; on the latter dg vector space, we define a bicoderivation *via*

$$(1) \quad d_{K_1} \otimes 1 \otimes 1 + 1 \otimes 1 \otimes d_{K_2} - 1 \otimes 1 \otimes d_B \otimes 1 \otimes 1,$$

where 1 denotes here the identity operator on the corresponding factor.

Remark 3.1. We now need a *caveat* regarding the fact that we use suspension and de-suspension in order to construct an A_{∞} -structure on the tensor product of two A_{∞} -bimodules, namely, while suspension or de-suspension of an A_{∞} -(bi)module is again an A_{∞} -(bi)module in a natural way, this is not true anymore for an A_{∞} -algebra. In fact, an A_{∞} -algebra structure on A cannot be transported on $A[-1]$, because of the fact that twisting w.r.t. suspension and de-suspension of the Taylor components of the A_{∞} -structure on A would produce maps of the wrong degree.

It is more instructive to write down explicit formulæ for the Taylor components of the previous bicoderivation

$$\begin{aligned}
(2) \quad d_{K_1 \otimes_B K_2}^{m,n}(a_1 | \cdots | a_m | k_1 \otimes (b_1 | \cdots | b_q) \otimes k_2 | c_1 | \cdots | c_n) &= 0, \quad m, n > 0 \\
d_{K_1 \otimes_B K_2}^{m,0}(a_1 | \cdots | a_m | k_1 \otimes (b_1 | \cdots | b_q) \otimes k_2) &= \sum_{l=0}^q s \left(s^{-1} (d_{K_1}^{m,l}(a_1 | \cdots | a_m | k_1 | b_1 | \cdots | b_l)) \otimes (b_{l+1} | \cdots | b_q) \otimes k_2 \right), \quad m > 0 \\
d_{K_1 \otimes_B K_2}^{0,n}(k_1 \otimes (b_1 | \cdots | b_q) \otimes k_2 | c_1 | \cdots | c_n) &= (-1)^{|k_1| + \sum_{j=1}^q (|b_j| - 1)} \sum_{l=0}^q s (k_1 \otimes (b_1 | \cdots | b_l) \otimes \\
&\quad s^{-1} (d_{K_2}^{q-l,n}(b_{l+1} | \cdots | b_q | k_2 | c_1 | \cdots | c_n))), \quad n > 0, \\
d_{K_1 \otimes_B K_2}^{0,0}(s(k_1 \otimes (b_1 | \cdots | b_q) \otimes k_2)) &= \sum_{l=0}^q s \left(s^{-1} (d_{K_2}^{0,l}(k_1 | b_1 | \cdots | b_l) \otimes (b_{l+1} | \cdots | b_q) \otimes k_2) \right) + \\
&\quad + \sum_{\substack{0 \leq l \leq q \\ 0 \leq p \leq q-l}} (-1)^{(|k_1| - 1) + \sum_{j=1}^l (|b_j| - 1)} s(k_1 \otimes (b_1 | \cdots | b_l | d_B^p(b_{l+1} | \cdots | b_{l+p}) | \cdots | b_q) \otimes k_2) + \\
&\quad + (-1)^{|k_1| + \sum_{j=1}^q (|b_j| - 1)} \sum_{l=0}^q s \left(k_1 \otimes (b_1 | \cdots | b_l) \otimes s^{-1} (d_{K_2}^{q-l,0}(b_{l+1} | \cdots | b_q | k_2)) \right).
\end{aligned}$$

Remark 3.2. We observe that the signs in (2) are dictated by Koszul's sign rule, together with the signs arising from the previous isomorphism $T(A[1]) \otimes (K_1 \otimes_B K_2)[1] \otimes T(C[1]) \cong T(A[1]) \otimes K_1[1] \otimes T(B[1]) \otimes K_2[1] \otimes T(C[1])$ of said degree -1 .

Proposition 3.3. *For A_∞ -algebras A, B, C and A_∞ -bimodules K_1, K_2 as above, $(K_1 \otimes_B K_2, d_{K_1 \otimes_B K_2})$, where the bicoderivation $d_{K_1 \otimes_B K_2}$ is defined in (2), is an A_∞ - A - C -bimodule.*

The proof of Proposition 3.3 follows from the following argument: since the bicoderivation $d_{K_1 \otimes_B K_2}$ is a twist of (1), it suffices to prove that (1) squares to 0. Using the fact that d_B, d_{K_1} and d_{K_2} all square to 0, we are reduced to prove that

$$\begin{aligned}
(3) \quad & (d_{K_1} \otimes 1 \otimes 1)(1 \otimes 1 \otimes d_{K_2}) - (d_{K_1} \otimes 1 \otimes 1)(1 \otimes 1 \otimes d_B \otimes 1 \otimes 1) + \\
& (1 \otimes 1 \otimes d_{K_2})(d_{K_1} \otimes 1 \otimes 1) - (1 \otimes 1 \otimes d_{K_2})(1 \otimes 1 \otimes d_B \otimes 1 \otimes 1) - \\
& - (1 \otimes 1 \otimes d_B \otimes 1 \otimes 1)(d_{K_1} \otimes 1 \otimes 1 + 1 \otimes 1 \otimes d_{K_2})
\end{aligned}$$

vanishes.

Since d_{K_1} and d_{K_2} are bicoderivations, and d_B is a coderivation, the sum of the first and second line in (3) can be rewritten as

$$\begin{aligned}
& 1 \otimes 1 \otimes d_B \otimes 1 \otimes d_C + (1 \otimes 1 \otimes d_B \otimes (\text{pr}_{K_2} \circ d_{K_2}))(1 \otimes 1 \otimes \Delta_B \otimes 1 \otimes 1) - \\
& - d_A \otimes 1 \otimes d_B \otimes 1 \otimes d_A - ((\text{pr}_{K_1} \circ d_{K_1}) \otimes d_B \otimes 1 \otimes 1)(1 \otimes 1 \otimes \Delta_B \otimes 1 \otimes 1),
\end{aligned}$$

where we have used the short-hand notation

$$(4) \quad \text{pr}_{K_1} \circ d_{K_1} = (1 \otimes (\text{pr}_{K_1} \circ d_{K_1}) \otimes 1)(\Delta_A \otimes 1 \otimes 1),$$

pr_{K_1} denoting the projection from $T(A[1]) \otimes K_1[1] \otimes T(B[1])$ onto $K_1[1]$ and *e.g.* Δ_A is the comultiplication of $T(A[1])$; similar notation holds for $\text{pr}_{K_2} \circ d_{K_2}$. The notation $\text{pr}_{K_1} \circ d_{K_1}$ means therefore simply that we select only that part of d_{K_1} given by the Taylor components $d_{K_1}^{m,n}$, thus forgetting about the action of d_B and d_C on the left and on the right, and similarly for $(\text{pr}_{K_1} \circ d_{K_1})$.

Using the same arguments, together with the nilpotence of d_B , yields the same expression for the last line of (3), up to global minus sign, coming from Koszul's sign rule.

Remark 3.4. We point out that, in general, the right-hand side of (4) should be

$$(1 \otimes (\text{pr}_{K_1} \circ d_{K_1}) \otimes 1)(\Delta_A \otimes 1 \otimes \Delta_B),$$

but, in this framework, the second coproduct Δ_B would be redundant and would produce too many terms, due to the fact that there is already a coproduct Δ_B , whence the reason, why we suppressed it in (4).

Alternatively, straightforward computations involving quadratic relations w.r.t. the Taylor components of d_A, d_C and $d_{K_1 \otimes_B K_2}$, which in turn can be re-written (after a very tedious book-keeping of all signs involved) in terms of the quadratic relations between the Taylor components of the codifferentials on A, B, C, K_1 and K_2 . Such computations are similar to the computations in [8, Chapter 4] in the case of right A_∞ -modules.

It is easy to verify that, if both A, C are flat, then the Taylor component $m_{K_1 \underline{\otimes}_B K_2}^{0,0}$ yields a structure of dg vector space on $K_1 \underline{\otimes}_B K_2$ (while B may be curved).

3.1. The A_∞ -bar construction of an A_∞ -bimodule. We consider two (possibly curved) A_∞ -algebras, and an A_∞ - A - B -bimodule K . It is easy to verify that e.g. A can be endowed with the structure of an A_∞ - A - A -bimodule, whose Taylor components are specified *via* the assignment $d_A^{m,n} = d_A^{m+1+n}$.

In particular, we may form the tensor product $A \underline{\otimes}_A K$, which has the structure of an A_∞ - A - B -bimodule, according to Proposition 3.3; similarly, we may consider the A_∞ - A - B -bimodule $K \underline{\otimes}_B B$.

An A_∞ -algebra A is said to be **unital**, if it possesses an element 1 of degree 0, such that

$$m_A^2(1 \otimes a) = m_A^2(a \otimes 1) = a, \quad m_A^n(a_1 \otimes \cdots \otimes a_n) = 0, \quad n \neq 2,$$

if $a_i = 1$, for some $i = 1, \dots, n$. If A is unital, and K is an A_∞ - A - B -bimodule, then K is (left-)unital w.r.t. A , if the identities hold true

$$m_K^{1,0}(1 \otimes k) = k, \quad m_K^{m,n}(a_1 \otimes \cdots \otimes a_m \otimes k \otimes b_1 \otimes \cdots \otimes b_n) = 0, \quad m \neq 1, \quad n \geq 0,$$

if $a_i = 1$, for some $i = 1, \dots, m$.

Now, there is a natural morphism μ of A_∞ - A - B -bimodules from $A \underline{\otimes}_A K$ to K : the cofreeness of A_∞ - A - B -bimodules implies that such a morphism is uniquely specified by its Taylor components

$$\mu^{m,n} : A[1]^{\otimes m} \otimes (A \otimes A[1]^{\otimes q} \otimes K)[1] \otimes B[1]^{\otimes n} \cong A[1]^{\otimes m+1+q} \otimes K[1] \otimes B[1]^{\otimes n} \rightarrow K[1], \quad m, n, q \geq 0,$$

all of degree 0, which have to additionally satisfy an infinite family of quadratic relations involving the Taylor components of the A_∞ -structures on $A \underline{\otimes}_A K$ and K . We notice that the isomorphism in the middle is of degree -1 .

There is a natural candidate for the morphism μ , namely,

$$(5) \quad \begin{aligned} & \mu^{m,n}(a_1 | \cdots | a_m | a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes k | b_1 | \cdots | b_n) = \\ & = (-1)^{\sum_{i=1}^m (|a_i| - 1) + |a| + \sum_{j=1}^q (|\tilde{a}_j| - 1)} d_K^{m+1+q,n}(a_1 | \cdots | a_m | a | \tilde{a}_1 | \cdots | \tilde{a}_q | k | b_1 | \cdots | b_n), \quad m, n, q \geq 0. \end{aligned}$$

Similar formulæ hold true for the case of the A_∞ - A - B -bimodule $K \underline{\otimes}_B B$.

Proposition 3.5. *For two (possibly curved) A_∞ -algebras A, B and an A_∞ - A - B -bimodule K , there is a natural morphism μ , defined by (5), of A_∞ - A - B -bimodules from $A \underline{\otimes}_A K$ to K .*

If A, B are both flat, and A, K are (left-)unital, then the A_∞ -morphism (5) is an A_∞ -quasi-isomorphism.

We first observe that the Taylor components in (5) have the right degree. Then, the proof of Proposition 3.5 consists in checking the aforementioned quadratic identities: in view of (2), and after a straightforward, but tedious, book-keeping of all signs involved, such quadratic identities can be re-written as the quadratic identities for the Taylor components of the A_∞ -structures on A and K , whence the first claim follows.

If A, B are both flat, then $d_{A \underline{\otimes}_A K}^{0,0}$ endows $A \underline{\otimes}_A K$ with the structure of a dg vector space: μ is an A_∞ -quasi-isomorphism, if and only if its Taylor component $\mu^{0,0}$ (which is automatically a morphism of dg vector spaces) induces an isomorphism in cohomology.

If A and K are unital, we set

$$(6) \quad \nu^{0,0}(k) = 1 \otimes k,$$

$$(7) \quad \sigma(a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes k) = 1 \otimes (a | \tilde{a}_1 | \cdots | \tilde{a}_q) \otimes k, \quad q \geq 0.$$

It is not difficult to check that the degree of (6) is 0, while the degree of (7) is -1 . Moreover, direct computations involving the fourth identity in (2) and the fact that A and K are unital imply that (6) is homotopically inverse to $\mu^{0,0}$, with explicit homotopy (7), whence the second claim follows.

Remark 3.6. We observe that Proposition 3.5 has been stated and proved in the framework of right A_∞ -modules in [8].

3.2. The A_∞ -bimodule structure on the bar resolution of the augmentation module. For V as in Section 2, we consider the symmetric algebra $A = S(V^*)$ and the exterior algebra $B = \wedge(V)$: both are unital dg algebras with trivial differential, A is concentrated in degree 0, while B is non-negatively graded. In particular, A and B can be viewed as flat, unital A_∞ -algebras, whose only non-trivial Taylor components are d_A^2 and d_B^2 respectively.

According to [1], $K = \mathbb{K}$ can be endowed with a non-trivial A_∞ - A - B -bimodule structure, which restricts to the natural augmentation left- and right-modules; non-triviality, here, means that there are non-trivial Taylor components $d_K^{m,n}$, for both m and n non-zero, e.g.

$$m_K^{1,1}(a \otimes 1 \otimes b) = \langle b, a \rangle, \quad a \in V, \quad b \in V^*,$$

and $\langle \bullet, \bullet \rangle$ denotes the duality pairing between V^* and V . For a complete description of the A_∞ - A - B -bimodule structure on K , we refer to [1, Subsection 6.2].

Since A, B are flat, and A, K are unital, Proposition 3.5 implies that there is an A_∞ -quasi-isomorphism of A_∞ - A - B -bimodules from $A \underline{\otimes}_A K$ to K . A direct computation implies that $A \underline{\otimes}_A K$ is a dg vector space concentrated in non-positive degrees; recalling (2), its A_∞ - A - B -bimodule structure is given by

$$(8) \quad \begin{aligned} d_{A \underline{\otimes}_A K}^{0,0}(a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes 1) &= s \left((a \tilde{a}_1) \otimes (\tilde{a}_2 | \cdots | \tilde{a}_q) \otimes 1 + \sum_{i=1}^{q-1} (-1)^i a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_i \tilde{a}_{i+1} | \cdots | \tilde{a}_q) \otimes 1 + \right. \\ &\quad \left. + (-1)^q a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_{q-1}) \otimes \tilde{a}_q(0) \right), \\ d_{A \underline{\otimes}_A K}^{1,0}(a_1 | a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes 1) &= s((a a_1) \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes 1), \\ d_{A \underline{\otimes}_A K}^{0,n}(a \otimes (\tilde{a}_1 | \cdots | \tilde{a}_q) \otimes 1 | b_1 | \cdots | b_n) &= (-1)^q \sum_{l=0}^q s \left(a_1 \otimes (\tilde{a}_1 | \cdots | \tilde{a}_l) \otimes s^{-1}(d_K^{q-l,n}(\tilde{a}_{l+1} | \cdots | \tilde{a}_q | 1 | b_1 | \cdots | b_n)) \right), \end{aligned}$$

and in all other cases, the Taylor components are trivial.

The first identity in (8) yields the identification between the dg vector space $(A \underline{\otimes}_A K, d_{A \underline{\otimes}_A K}^{0,0})$ identifies with the bar complex of the left augmentation module $K = \mathbb{K}$ over A .

Further, the identities (8) imply that the left A_∞ - A -module structure on the bar complex $A \underline{\otimes}_A K$ of K is the standard one, while the non-triviality of the A_∞ - A - B -bimodule structure on K yields non-triviality of the right A_∞ - B -module structure on $A \underline{\otimes}_A K$.

4. THE KOSZUL COMPLEX OF $A = S(V^*)$: A BRIEF *memento*

For V as in Section 2, we consider the symmetric algebra $A = S(V^*)$ and the exterior algebra $B = \wedge(V)$.

We further consider the Koszul complex $K(V)$ of A . As a graded vector space,

$$K^q(V) = \wedge_{\mathcal{O}_V}^{-q} \Omega_{\mathcal{O}_V/\mathbb{K}}^1, \quad q \geq 0,$$

where $\Omega_{\mathcal{O}_V/\mathbb{K}}^1$ denotes the module of Kaehler differentials on $\mathcal{O}_V = S(V^*) = A$ as a $\mathbb{K}$ -algebra; in particular, $K(V)$ is non-positively graded. The differential ∂ on $K(V)$ is induced by the (left) contraction w.r.t. the Euler vector field on V ; further, $K(V)$ admits an obvious left A -action, and contraction w.r.t. polyvector fields on V induces by restriction (and keeping track of Koszul's sign rule) a right B -action on $K(V)$. In particular, $K(V)$ admits the structure of an A_∞ - A - B -bimodule, whose only non-trivial Taylor components are labeled by pairs of indices (m, n) , such that $m + n \leq 1$. For later computations, we choose a $\mathbb{K}$ -basis $\{x_i\}$ on V^* as in Section 2: in particular, we may write

$$K(V) \cong \mathbb{K}[x_i, \theta_j],$$

where $\{\theta_j\}$ denotes a set of odd coordinates of degree -1 , which anticommute with each other and commute with x_i ; w.r.t. the previous algebra isomorphism, $x_i \mapsto x_i$, $dx_i \mapsto \theta_i$, and ∂ is uniquely determined by the graded Leibniz rule (from the left) and by $\partial(x_i) = 0$, $\partial(\theta_i) = x_i$.

Furthermore, we may also write $B = \mathbb{K}[\partial_{\theta_j}]$, where the partial derivative ∂_{θ_j} has degree 1, and acts in an obvious way from the left on $K(V)$: thus, the right B -action on $K(V)$ takes the explicit form

$$K(V) \otimes B \ni \eta \otimes b_1 \mapsto (-1)^{|\eta||b_1|} b_1(\eta).$$

We finally observe that $(K(V), \partial)$ is a free resolution of the left A -module K *via* augmentation; it is also a free resolution of the right B -module K *via* augmentation, because of the isomorphism $B \cong (\mathbb{K}[\theta_j])[-d]$ induced by contraction w.r.t. polyvector fields.

5. AN EXPLICIT A_∞ -QUASI-ISOMORPHISM BETWEEN THE BAR AND THE KOSZUL COMPLEX OF A

We use the same notation as in the preceding sections; we only observe that in the whole Section, $A = S(V^*)$, $B = \wedge(V)$ and $K = \mathbb{K}$ with the A_∞ - A - B -bimodule structure from [1].

Since $A \underline{\otimes}_A K$ and $K(V)$ are both resolutions of the left augmentation module $K = \mathbb{K}$ over A , arguments from abstract (co)homological algebra imply that they are quasi-isomorphic to each other as complexes of free left A -modules. More precisely, the quasi-isomorphism from $K(V)$ to $A \underline{\otimes}_A K$ as complexes of left A -modules has an explicit form, namely

$$(9) \quad \Phi(\theta_{i_1} \cdots \theta_{i_q}) = \sum_{\sigma \in \mathfrak{S}_q} (-1)^\sigma 1 \otimes (x_{\sigma(i_1)} | \cdots | x_{\sigma(i_q)}) \otimes 1, \quad 1 \leq i_1 < \cdots < i_q \leq d.$$

It suffices to define the morphism Φ on monomials of the form $\theta_{i_1} \cdots \theta_{i_q}$, and then extend it A -linearly on the left. It follows immediately that Φ is of degree 0 and commutes with left A -action. An easy computation shows that Φ commutes with differentials. It is a quasi-isomorphism since $\Phi(1) = 1$.

Theorem 5.1. *For a finite-dimensional $\mathbb{K}$ -vector space V , the morphism (9) extends to a quasi-isomorphism of A_∞ - A - B -bimodules from $K(V)$ to $A \underline{\otimes}_A K$, where the A_∞ - A - B -bimodule structures on $K(V)$ and $A \underline{\otimes}_A K$ are described in Sections 4 and 3.1 respectively.*

Proof. We know that (9) is a morphism of degree 0 from $K(V)$ to $A \underline{\otimes}_A K$: we declare (the conjugation w.r.t. s of) Φ to be the $(0, 0)$ -th Taylor component of the desired A_∞ -quasi-isomorphism, while for (m, n) such that $m + n \geq 1$, we set simply 0.

By the previous arguments, the only non-trivial identities to check are

$$(10) \quad d_{A \underline{\otimes}_A K}^{0,1}(\Phi(\eta)|b_1) = \Phi(d_{K(V)}^{0,1}(\eta|b_1)),$$

$$(11) \quad d_{A \underline{\otimes}_A K}^{0,n}(\Phi(\eta)|b_1| \cdots |b_n) = 0, \quad n \geq 2,$$

for b_i , $i = 1, \dots, n$, resp. η , a general element of B , resp. $K(V)$.

We begin by proving Identity (11): A -linearity implies that we may take η of the form $\theta_{i_1} \cdots \theta_{i_q}$, $1 \leq i_1 < \cdots < i_q \leq d$. Recalling now Identity (9), we rewrite the left-hand side in (11) as

$$d_{A \underline{\otimes}_A K}^{0,n}(\Phi(\eta)|b_1| \cdots |b_n) = (-1)^q \sum_{l=0}^q \sum_{\sigma \in \mathfrak{S}_q} (-1)^\sigma s \left(1 \otimes (x_{\sigma(i_1)} | \cdots | x_{\sigma(i_l)}) \otimes s^{-1}(d_K^{q-l,n}(x_{\sigma(i_{l+1}}) | \cdots | x_{\sigma(i_q)} | 1 | b_1 | \cdots | b_n) \right).$$

We now analyze the last factor on the right-hand side: degree reasons imply that, for $0 \leq l \leq q$,

$$-(q-l) - 1 + \sum_{j=1}^n (|b_j| - 1) + 1 \stackrel{!}{=} -1 \iff \sum_{j=1}^n |b_j| = n + q - l - 1 > q - l,$$

because $n \geq 2$ by assumption.

We recall now from [1, Subsection 6.2] that the Taylor components of the A_∞ - A - B -bimodule structure on K are constructed explicitly *via* admissible graphs, in a way reminiscent of Kontsevich's graphical technique of [7]: using the same notation of [7], admissible graphs of type (m, n) (elements of $\mathcal{G}_{m,n}$) are graphs embedded in $\mathbb{R} \sqcup \mathbb{H}$ with m vertices of the first type (*i.e.* lying in the complex upper half-plane $\mathbb{H}$), n vertices of the second type (*i.e.* n ordered vertices on the real axis $\mathbb{R}$), and with a certain number of oriented edges between them. In the present framework, we consider typically elements of $\mathcal{G}_{0,m+1+n}$, where to the first m vertices of the second type we associate elements of A , to the $m+1$ -st vertex of the second type 1 as an element of $\mathbb{K}$, and to the last n vertices of the second type we associate elements of B : accordingly, oriented edges are associated to elements of B , which we view as translation-invariant poly-derivations acting on $A = \mathcal{O}_V$. To such admissible graphs are associated polydifferential operators on A , B and K with values in K (by the obvious rule that oriented edges correspond to derivatives) and integral weights, for whose precise treatment we refer to [1, Subsection 6.2] again: suffice it to recall here that the integral weight of a given admissible graph Γ in $\mathcal{G}_{0,m+1+n}$ is the integral over the compactified configuration space $\mathcal{C}_{0,m+1+n}^+$ of $m+1+n$ ordered points on the real axis (modulo rescalings and real translations) of a differential form depending explicitly on Γ , roughly defined *via* the rule that a closed 1-form is associated to an edge connecting two vertices.

The previous strict inequality, which is a consequence of the non-vanishing of the integral weight of any admissible graph appearing in the formula for $d_K^{q-l,n}$, implies the claim: in fact, any admissible graph Γ in $\mathcal{G}_{0,n+q-l-1}$ in $d_K^{q-l,n}$ must have exactly $n + q - l - 1$ arrows departing from the vertices on the right-hand side of the $q - l + 1$ -st vertex and incoming on the vertices on the left-hand side; no other arrows or loops are allowed. Since $n \geq 2$ and A , B and K are unital, then all vertices (except the $q - l + 1$ -st vertex) must be at least univalent: more precisely, the vertices on the left-hand side of the $q - l + 1$ -st vertex must have exactly one incoming arrow (because of degree reasons), while the vertices on the right-hand side must have at least one outgoing arrow, and the previous strict inequality proves that no such graphs exist.

It remains to prove Identity (10). We first evaluate the right-hand side: using the isomorphism at the end of Section 4, we may write $b_1 = \partial_{\theta_{j_1}} \cdots \partial_{\theta_{j_p}}$, whence the right-hand side takes the form $(-1)^{(p+1)q} \Phi(b_1(\eta))$.

The left-hand side of Identity (10) has the explicit form

$$\begin{aligned} d_{A \otimes_A K}^{0,1}(\Phi(\eta)|b_1) &= (-1)^q \sum_{l=0}^q \sum_{\sigma \in \mathfrak{S}_q} (-1)^\sigma s \left(1 \otimes (x_{\sigma(i_1)} | \cdots | x_{\sigma(i_l)}) \otimes s^{-1}(d_K^{q-l,1}(x_{\sigma(i_{l+1}}) | \cdots | x_{\sigma(i_q)} | 1 | b_1)) \right) = \\ &= (-1)^q \sum_{\sigma \in \mathfrak{S}_q} (-1)^\sigma s \left(1 \otimes (x_{\sigma(i_1)} | \cdots | x_{\sigma(i_{q-p})}) \otimes s^{-1}(d_K^{p,1}(x_{\sigma(i_{q-p+1}}) | \cdots | x_{\sigma(i_q)} | 1 | b_1)) \right), \end{aligned}$$

where the second equality follows because of degree reasons.

First, if $q \leq p-1$, both sides of Identity (10) vanish: this follows immediately from the previous formulæ. It remains therefore to prove the claim in the case $q \leq p$.

We now take a closer look at the left-hand side of Identity (10): we need to understand, in this particular case, the Taylor component $d_K^{p,1}$. We assume $p \geq 1$, because the case $p = 0$ follows immediately by direct computations and previous considerations. In view of [1, Subsection 6.2], $d_K^{p,1}$ is a sum over admissible graphs Γ in $\mathcal{G}_{0,p+2}$: in fact, there is only one such admissible graph contributing non-trivially, pictorially




Figure 1 - The only admissible graph contributing non-trivially to $d_K^{p,1}(x_{\sigma(i_{q-p+1})} | \cdots | x_{\sigma(i_q)} | 1 | b_1)$

The differential operator $\mathcal{O}_\Gamma$ associated to the admissible graph Γ as in Figure 1 can be explicitly evaluated, following the prescriptions in [1, Subsection 6.2], namely

$$(12) \quad s^{-1}(d_K^{p,1}(x_{\sigma(i_{q-p+1})} | \cdots | x_{\sigma(i_q)} | 1 | b_1)) = \frac{(-1)^p}{p!} b_1 (\theta_{\sigma(i_{q-p+1})} \cdots \theta_{\sigma(i_q)}),$$

The claim now follows from Identity (12). □

5.1. A final remark. In this final Subsection, we want to point out that the techniques portrayed in Sections 3, 4 and in the present one apply as well to the more general situation examined in [1]: namely, for any two subspaces U_i , $i = 1, 2$, of a finite-dimensional $\mathbb{K}$ -vector space V , we may associate an A_∞ -category with two objects, *i.e.* U_1 and U_2 , two A_∞ -algebras A and B and an A_∞ - A - B -bimodule K .

More explicitly, A , resp. B , is the A_∞ -algebra associated to the graded vector space of global (regular) sections of the exterior algebra of the normal bundle of U_1 , resp. of U_2 , in V (with obvious product and trivial differential); K is the graded vector space of global (regular) sections of the exterior algebra of the quotient bundle $TV/(TU_1 + TU_2)$ over $U_1 \cap U_2$, and the A_∞ -bimodule structure extends the natural left A - and right B -action: of course, A , resp. B , resp. K , must be understood as the endomorphism space of U_1 , resp. of U_2 , resp. the space of morphisms from U_1 to U_2 , declaring trivial the remaining one. We do not indulge in its explicit construction: suffice it to say that it involves (once again) Kontsevich's diagrammatic techniques, and we observe that it reduces, when $U_1 = V$ and $U_2 = \{0\}$, to the one we have made explicit in the previous computations.

As already remarked in [1, Subsection 7.3], there is also a Koszul complex $K(V, U_1, U_2)$ *e.g.* for A , whose construction is similar to the one of $K(V)$ in Section 4, with obvious due changes; again, $K(V, U_1, U_2)$ can be given the structure of a DG A - B -bimodule (hence, of an A_∞ - A - B -bimodule).

Furthermore, there is a natural (graded) version of the quasi-isomorphism (9) from the Koszul complex $K(V, U_1, U_2)$ to the bar resolution of K as a left A -module: then, a more involved vanishing lemma, proved in the same spirit of Theorem 5.1, Identity (10), and a more refined version of Identity (11), imply that such a quasi-isomorphism extends to an A_∞ -quasi-isomorphism of A - B -bimodules. In particular, Theorem 1.1 holds true in the more general situation of [1].

Since Keller's condition in [10] holds true in the more general situation already examined, *i.e.* for a DG category with two objects U_i , $i = 1, 2$, and with spaces of morphisms given by A , B (the endomorphisms of U_1 and U_2 respectively) and $K(V, U_1, U_2)$ (the morphisms from U_1 to U_2), a natural question is, if it is possible to formulate and prove the main result of [10] in this more general setting, using Tamarkin's results instead of Kontsevich's.

REFERENCES

- [1] Damien Calaque, Giovanni Felder, Andrea Ferrario, and Carlo A. Rossi, *Bimodules and branes in deformation quantization* (2009), available at [arXiv:0908.2299](https://arxiv.org/abs/0908.2299).
- [2] Damien Calaque, Giovanni Felder, and Carlo A. Rossi, *Deformation quantization with generators and relations* (2009), available at [arXiv:0911.4377](https://arxiv.org/abs/0911.4377).
- [3] Alberto S. Cattaneo and Giovanni Felder, *Coisotropic submanifolds in Poisson geometry and branes in the Poisson sigma model*, Lett. Math. Phys. **69** (2004), 157–175. MR2104442 (2005m:81285)
- [4] ———, *Relative formality theorem and quantisation of coisotropic submanifolds*, Adv. Math. **208** (2007), no. 2, 521–548. MR2304327 (2008b:53119)
- [5] Bernhard Keller, *Introduction to A-infinity algebras and modules*, Homology Homotopy Appl. **3** (2001), no. 1, 1–35 (electronic). MR1854636 (2004a:18008a)
- [6] ———, *Derived invariance of higher structures on the Hochschild complex* (2003), available at <http://people.math.jussieu.fr/~keller/publ/dih.pdf>.
- [7] Maxim Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216. MR2062626 (2005i:53122)
- [8] Kenji Lefèvre-Hasegawa, *Sur les A_∞ -catégories* (2003), available at <http://people.math.jussieu.fr/~keller/lefevre/TheseFinale/tel-00007761>.
- [9] Boris Shoikhet, *Kontsevich formality and PBW algebras* (2007), available at [arXiv:0708.1634](https://arxiv.org/abs/0708.1634).
- [10] ———, *Koszul duality in deformation quantization and Tamarkin's approach to Kontsevich formality*, to appear in Adv. Math. (2008), available at [arXiv:0805.0174](https://arxiv.org/abs/0805.0174).
- [11] Thomas Willwacher, *A counterexample to the quantizability of modules*, Lett. Math. Phys. **81** (2007), no. 3, 265–280. MR2355492 (2008j:53160)

A. F. AND T. W.: DEPARTMENT OF MATHEMATICS, ETH ZURICH, 8092 ZURICH, SWITZERLAND

C. A. R.: CENTRO DE ANÁLISE MATEMÁTICA, GEOMETRIA E SISTEMAS DINÂMICOS, DEPARTAMENTO DE MATEMÁTICA, INSTITUTO SUPERIOR TÉCNICO, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL